

(b) The pharmacological and toxicological activity of the diagnostic radiopharmaceutical (including any carrier or ligand component of the diagnostic radiopharmaceutical), and

(c) The estimated absorbed radiation dose of the diagnostic radiopharmaceutical.

#### § 315.4 Indications.

(a) For diagnostic radiopharmaceuticals, the categories of proposed indications for use include, but are not limited to, the following:

- (1) Structure delineation;
- (2) Functional, physiological, or biochemical assessment;
- (3) Disease or pathology detection or assessment; and
- (4) Diagnostic or therapeutic patient management.

(b) Where a diagnostic radiopharmaceutical is not intended to provide disease-specific information, the proposed indications for use may refer to a biochemical, physiological, anatomical, or pathological process or to more than one disease or condition.

#### § 315.5 Evaluation of effectiveness.

(a) The effectiveness of a diagnostic radiopharmaceutical is assessed by evaluating its ability to provide useful clinical information related to its proposed indications for use. The method of this evaluation varies depending upon the proposed indication(s) and may use one or more of the following criteria:

- (1) The claim of structure delineation is established by demonstrating in a defined clinical setting the ability to locate anatomical structures and to characterize their anatomy.
- (2) The claim of functional, physiological, or biochemical assessment is established by demonstrating in a defined clinical setting reliable measurement of function(s) or physiological, biochemical, or molecular process(es).
- (3) The claim of disease or pathology detection or assessment is established by demonstrating in a defined clinical setting that the diagnostic radiopharmaceutical has sufficient accuracy in identifying or characterizing the disease or pathology.
- (4) The claim of diagnostic or therapeutic patient management is estab-

lished by demonstrating in a defined clinical setting that the test is useful in diagnostic or therapeutic patient management.

(5) For a claim that does not fall within the indication categories identified in § 315.4, the applicant or sponsor should consult FDA on how to establish the effectiveness of the diagnostic radiopharmaceutical for the claim.

(b) The accuracy and usefulness of the diagnostic information is determined by comparison with a reliable assessment of actual clinical status. A reliable assessment of actual clinical status may be provided by a diagnostic standard or standards of demonstrated accuracy. In the absence of such diagnostic standard(s), the actual clinical status must be established in another manner, e.g., patient followup.

#### § 315.6 Evaluation of safety.

(a) Factors considered in the safety assessment of a diagnostic radiopharmaceutical include, among others, the following:

- (1) The radiation dose;
- (2) The pharmacology and toxicology of the radiopharmaceutical, including any radionuclide, carrier, or ligand;
- (3) The risks of an incorrect diagnostic determination;
- (4) The adverse reaction profile of the drug;
- (5) Results of human experience with the radiopharmaceutical for other uses; and
- (6) Results of any previous human experience with the carrier or ligand of the radiopharmaceutical when the same chemical entity as the carrier or ligand has been used in a previously studied product.

(b) The assessment of the adverse reaction profile includes, but is not limited to, an evaluation of the potential of the diagnostic radiopharmaceutical, including the carrier or ligand, to elicit the following:

- (1) Allergic or hypersensitivity responses,
- (2) Immunologic responses,
- (3) Changes in the physiologic or biochemical function of the target and nontarget tissues, and
- (4) Clinically detectable signs or symptoms.